JOURNAL OF CHROMATOGRAPHY A

VOL. 773, NOS. 1 + 2 27 JUNE 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Pressure drop in centrifugal partition chromatography
by M.J. van Buel, L.A.M. van der Wielen and K.Ch.A.M. Luyben (Delft, The Netherlands) (Received 17 February 1997) . . . 1

Modelling gradient elution in centrifugal partition chromatography
by M.J. van Buel, L.A.M. van der Wielen and K.Ch.A.M. Luyben (Delft, The Netherlands) (Received 17 February 1997) . . . 13

Detection of the oxidative products of 3-hydroxykynurenine using high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–electron spin resonance spectrometry and high-performance liquid chromatography–electrochemical detection–ultraviolet absorption detection–mass spectrometry
by H. Iwahashi and T. Ishii (Wakayama, Japan) (Received 18 February 1997) 23

Study of the physico-chemical properties of some packing materials. III. Pore size and surface area distribution
by H. Guan-Sajonz and G. Guiochon (Knoxville, TN, USA and Oak Ridge, TN, USA) and E. Davis, K. Gulakowski and D.W. Smith (Norcross, GA, USA) (Received 31 January 1997) 33

Retention characteristics and selected applications of cyclic siloxane-based octadecylsilyl bonded phases in reversed-phase high-performance liquid chromatography
by S.O. Akapo (Moffett Field, CA, USA) and M.T. Matyska and J.J. Pesek (San Jose, CA, USA) (Received 18 February 1997) 53

Sol–gel-derived fluorinated stationary phase for open tubular electrochromatography
by P. Narang and L.A. Colón (Buffalo, NY, USA) (Received 21 January 1997) 65

Preparation and use of ion-exchange chromatographic supports based on perfluoropolymers
by G.E. McCreath, R.O. Owen, D.C. Nash and H.A. Chase (Cambridge, UK) (Received 17 January 1997) 73

Separation of enantiomers on a chiral stationary phase based on ovoglycoprotein. I. Influences of the pore size of base silica materials and bound protein amounts on chiral resolution
by J. Haginaka and H. Takehira (Nishinomiya, Japan) (Received 24 January 1997) 85

Reversed-phase separations of nitrogenous phospholipids on an octadecanoyl poly(vinyl alcohol) phase
by S.L. Abidi and T.L. Mounts (Peoria, IL, USA) (Received 24 February 1997) 93

Liquid chromatography on soft packing material, under axial compression. Size-exclusion chromatography of polypeptides
by A.V. Danilov, I.V. Vagenina, L.G. Mustaeva, S.A. Moshnikov, E.Yu. Gorbunova, V.V. Cherskii and M.B. Baru (Moscow, Russian Federation) (Received 6 January 1997) 103

Anion-exchange chromatography with electrochemical detection of alditols and sugars at a Cu_2O–carbon composite electrode
by T.R.I. Cataldi, D. Centonze and I.G. Casella (Potenza, Italy) and E. Desimoni (Milano, Italy) (Received 12 February 1997) 115

Determination of low-molecular-mass organic acids in fogwater by ion-exchange chromatography with automatic column switching
by Y. Suzuki (Kobe, Japan) (Received 23 January 1997) 123

Purification of chaperonins from thermophilic bacteria and archaea
by A. Joachimiak, E. Quaite-Randall and S. Tollaksen (Argonne, IL, USA), X. Mai and M.W.W. Adams (Athens, GA, USA), R. Josephs (Chicago, IL, USA) and C. Giometti (Argonne, IL, USA) (Received 3 February 1997) 131

Determination of phosphonates in natural waters by ion-pair high-performance liquid chromatography
by B. Nowack (St. Gallen, Switzerland) (Received 25 February 1997) 139

Evaluation of liquid chromatography methods for the analysis of benzylpenicillin and its related substances
by Z. Yongxin, A. Verhasselt and E. Roets (Leuven, Belgium), A. Perez and E. Porqueras (Madrid, Spain) and J. Hoogmartens (Leuven, Belgium) (Received 3 March 1997) 147

Speciation of chromium dyes by high-performance liquid chromatography with inductively coupled plasma mass spectrometric detection
by G.K. Zoorob and J.A. Caruso (Cincinnati, OH, USA) (Received 27 August 1996) 157

Determination of aspartame and phenylalanine in diet soft drinks by high-performance liquid chromatography with direct spectrofluorimetric detection
by K. Wróbel and K. Wróbel (Guanajuato, Mexico) (Received 21 February 1997) 163

New sorbents for removal of heavy metal ions: diamine-glow-discharge treated polyhydroxyethylmethacrylate microspheres
by A. Denizli, B. Salih and E. Pişkin (Ankara, Turkey) (Received 14 February 1997) 169

Analysis of colloids. IX. Investigation of the electrical double layer of colloidal inorganic nanometer-particles by size-exclusion chromatography
by Ch.-H. Fischer (Berlin, Germany) and E. Kenndler (Wien, Austria) (Received 26 February 1997) 179

Gas Chromatography

Semi-micro solid-phase extraction of organic compounds from aqueous and biological samples
by D.L. Mayer and J.S. Fritz (Ames, IA, USA) (Received 24 February 1997) 189

Synthesis and characterization of a novel wall coated capillary column for the separation of polycyclic aromatic hydrocarbons
by C.-Y. Liu, C.-C. Hu and C.-L. Yang (Taipei, Taiwan) (Received 17 February 1997) 199

Chemically bonded chelates as selective complexing sorbents for gas chromatography. VI. Modification of silica with $NiCl_2$ and $CoCl_2$ via β-diketonate groups
by W. Wasiak and I. Rykowska (Poznań, Poland) (Received 21 January 1997) 209

Capillary gas chromatography–mass spectrometry of lower oxyethylenated aliphatic alcohols
by K. Komárek, V. Pitthard and E. Kostrubaničová (Pardubice, Czech Republic), S. Škvarenina (Nováky, Slovakia) and J. Hoffmann (Zlín, Czech Republic) (Received 18 February 1997) 219

Absolute quantitation of lignin pyrolysis products using an internal standard
by P. Bocchini and G.C. Galletti (Bologna, Italy) and S. Camarero and A.T. Martinez (Madrid, Spain) (Received 24 January 1997) 227

Quantitative aspects of triglyceride analysis by gas–liquid chromatography using a short metal capillary column
by F. Ulberth and R. Gabernig (Vienna, Austria) (Received 27 February 1997) 233

Reaction of fatty acid amides and ethanolamides with trimethylsulfonium hydroxide
by K. Vosmann, E. Klein and N. Weber (Münster, Germany) (Received 31 January 1997) 239

Determination of amines in air and water using derivatization combined with solid-phase microextraction
by L. Pan, J.M. Chong and J. Pawliszyn (Waterloo, Canada) (Received 3 February 1997) 249

Chromatographic analysis of volatile sulphur compounds in wines using the static headspace technique with flame photometric detection
by M. Mestres, O. Busto and J. Guasch (Tarragona, Spain) (Received 14 February 1997) 261

Simultaneous quantitation of sixteen organochlorine pesticides in drinking waters using automated solid-phase extraction, high-volume injection, high-resolution gas chromatography
by W.C. Quayle, I. Jepson and I.A. Fowlis (Newcastle upon Tyne, UK) (Received 14 January 1997) 271

Supercritical Fluid Chromatography

Retention behavior of synthetic corticosteroids in packed-column supercritical fluid chromatography
by K. Yaku, K. Aoe, N. Nishimura and T. Sato (Osaka, Japan) and F. Morishita (Kyoto, Japan) (Received 21 February 1997) . 277

Planar Chromatography

Iodine–azide reagent in detection of thiophosphoryl nucleotides in thin-layer chromatography systems
by A. Kotyński, Z.H. Kudzin, A. Okruszek, D. Krajewska, M. Olesiak and A. Sierzchała (Łódź, Poland) (Received 19 November 1996) 285

Electrophoresis

Sensitive absorbance detection method for capillary electrophoresis based on laser wave-mixing
by Z. Wu and W.G. Tong (San Diego, CA, USA) (Received 4 February 1997) 291

Purification by reflux electrophoresis of whey proteins and of a recombinant protein expressed in *Dictyostelium discoideum*
by G.L. Corthals, B.M. Collins, B.C. Mabbutt, K.L. Williams and A.A. Gooley (Sydney, Australia) (Received 20 February 1997) 299

Enantiomeric separation of the novel growth hormone secretagogue MK-0677 by capillary zone electrophoresis
by L. Zhou, J. Trubig and A. Dovletoglou (Rahway, NJ, USA) and D.C. Locke (Flushing, NY, USA) (Received 27 January 1997) 311

Improved chiral separation using achiral modifiers in cyclodextrin modified capillary zone electrophoresis
by E. Billiot, J. Wang and I.M. Warner (Baton Rouge, LA, USA) (Received 21 February 1997) 321

Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis
by B. Chankvetadze, M. Saito, E. Yashima and Y. Okamoto (Nagoya, Japan) (Received 4 March 1997) 331

Development and validation of a capillary electrophoresis method within a pharmaceutical quality control environment and comparison with high-performance liquid chromatography
by G.S. Wynia, G. Windhorst, P.C. Post and F.A. Maris (Oss, The Netherlands) (Received 22 January 1997) 339

Capillary electrophoresis in metal analysis. Investigations of multi-elemental separation of metal chelates with aminopolycarboxylic acids
by A. Padarauskas (Vilnius, Lithuania) and G. Schwedt (Clausthal-Zellerfeld, Germany) (Received 16 January 1997) 351

SHORT COMMUNICATIONS

Column Liquid Chromatography

Quantitative determination of mono- and dipentaerythritol in synthesis solutions by high-performance liquid chromatography
by H. Li (Hunan, China) and Y. Li, J. Den and Q. Hu (Hunan, P.R. China) (Received 29 January 1997) 361

Separation of positional isomers of nitrobenzoic acid by reversed-phase liquid chromatography with 2-propanol–water–acetic acid as eluent
by P. Chen and M. Zhang (Beijing, China) (Received 7 February 1997) 365

Gas Chromatography

Influence of conformation on the accuracy of quantitative–structure retention relationship calculations in gas chromatography
by N. Dimov (Sofia, Bulgaria) and A. Osman (Bourgas, Bulgaria) (Received 30 December 1996) 368

Quantitative injections or calibrations in gas chromatography. Anomaly associated with syringe type
by C.R. French, C.J. Gray and R.S. Lehrle (Birmingham, UK) (Received 4 February 1997) 372

Supercritical Fluid Chromatography

Supercritical fluid chromatography–mass spectrometry of thyme extracts (*Thymus vulgaris* L.)
by C. Blum and K.-H. Kubeczka (Hamburg, Germany) and K. Becker (Bremen, Germany) (Received 28 February 1997) . . . 377

BOOK REVIEW 381

AUTHOR INDEX VOL. 773 383

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc